

Nilpotent Marsh and SUSY QM

V. P. Akulov

*City College of City University of New York,
138th St. Convent Ave , New York, NY 10031, USA*
and

Steven Duplij^{*†}

*Theory Group, Nuclear Physics Laboratory,
Kharkov State University, Kharkov 310077, Ukraine*

February 1, 2008

Abstract

We consider the nilpotent additions to classical trajectories in supersymmetric and nonsupersymmetric theories. The condition of anilpotence of action on some generalized solutions leads to the Witten supersymmetric Lagrangian. The condition of anilpotence of topological charge is the same as one of superpotential with spontaneous broken supersymmetry. We should vanish half of Grassmann constants of integration, because in this case only we obtain the same number of normalized bosonic and fermionic zero modes.

^{*}E-mail: Steven.A.Duplij@univer.kharkov.ua

[†]Internet: [http:// www-home.univer.kharkov.ua/~duplij](http://www-home.univer.kharkov.ua/~duplij)

1 Introduction

This paper is dedicated to Victor Isaakovich Ogievetsky who was an outstanding scientist and brilliant man. He gave a great contribution to quantum field theory, supersymmetry and supergravity.

In recent time the interest to extended supersymmetric quantum mechanics [1] with N supersymmetries [2, 3, 4, 5, 6, 7] is greatly renewed. This is connected on one side with dimensional reduction of additional dimensions of D-brane where one can receive SQM limit, which is more simple object. Maldacena's correspondence [8] between string theory on AdS spaces and superconformal quantum theories attracts attention to the structure of extremal black holes near the horizon which is described by the supersymmetric quantum mechanics [9].

On the other side the supermembrane in light cone gauge gives rise to SQM system with $SU(N)$ gauge symmetry [10, 5]. The D(-1) branes or D-instantons have the world volume which is just a point; they are topological objects and carry out topological charge.

Instantons as such solitons have zero modes connected with derivative by constants of integration. It is well known that these solutions (as a rule) break one half supersymmetries. For any supersymmetric theory the number of bosonic zero modes should be the same as fermionic ones.

We want to consider a full class of solutions for classical equations of motion, which consist of both bosonic and Grassmann constants of integration [11, 12, 13, 14, 4]. Fields take their values in Grassmann algebra with an even part and an odd part. But the action have to belong only to the even part and should not have any nilpotent addition, because the action is connected with a measure defined by functional integration in the path integral approach.

We here concentrate our attention mainly on the conditions which give absence of nilpotent additions to the action on the generalized classical trajectories and receive the connection between bosonic and fermionic parts of the interaction.

The plan of the paper is the following. In Section 2 we introduce the superfield approach to SQM with $N=2$ supersymmetries (to confine ourselves with most simple model). In Section 3 we consider full classical solutions and the corresponding nilpotent "marsh" — trajectories with nilpotent addition. In section 4 we investigate the condition of vanishing of half number of Grass-

mann constants, because only in this case we obtain exactly correct number of fermionic zero modes. In section 5 we obtain Witten's supersymmetric Lagrangian as a consequence of anilpotence of action on such trajectories. In section 6 we obtained SWKB rules for quantization. In section 7 we conclude with some comments.

2 Superfield Formulation

In $(1|2)$ dimensional Euclidean superspace and in the chiral basis $(\tau, \theta^+, \theta^-)$ a scalar superfield $\Phi(\tau, \theta^+, \theta^-)$ has the following expansion

$$\Phi(\tau, \theta^+, \theta^-) = q(\tau) + \theta^+ \psi_-(\tau) + \theta^- \psi_+(\tau) + \theta^+ \theta^- F(\tau), \quad (1)$$

where $q(\tau)$ is a bosonic coordinate, $F(\tau)$ is an auxiliary bosonic field, $\psi_{\pm}(\tau)$ are fermionic coordinates. Using chiral derivatives $D^{\pm} = \partial/\partial\theta^{\mp} + \theta^{\pm}\partial/\partial\tau$ the superfield Lagrangian of the model with a superpotential $V(\Phi)$ can be written as

$$\mathcal{L}_E = \frac{1}{2} D^+ \Phi \cdot D^- \Phi + V(\Phi) \quad (2)$$

and corresponding equations of motion are¹

$$\frac{1}{2} [D^+, D^-] \Phi - V'(\Phi) = 0. \quad (3)$$

The Euclidean action has the standard form

$$S_E = \int dt L_E, \quad (4)$$

where

$$L_E = \int d\theta^- d\theta^+ \mathcal{L}_E \quad (5)$$

is an ordinary Euclidean Lagrangian. Expanding (2) in θ -series which is finite due to nilpotence of $\theta^{\pm}$, using the nondynamical equation $F = -V'(q)$ and integrating by in (5) $\theta^{\pm}$ we obtain²

$$L_E = \frac{\dot{q}^2}{2} + \frac{V'^2(q)}{2} + \psi_+ \dot{\psi}_- + V''(q) \psi_+ \psi_-. \quad (6)$$

¹Prime is differentiation by argument.

²Dote is differentiation by τ .

3 Classical Solutions and Even Nilpotent Directions

From (6) we find the following equations of motion

$$\ddot{q} - V'(q) V''(q) + V'''(q) \psi_+ \psi_- = 0, \quad (7)$$

$$\dot{\psi}_\pm \mp V''(q) \psi_\pm = 0. \quad (8)$$

The classical solution of the system (7)–(8) is much simplified if one takes $\psi_\pm^{cl} = 0$ [15, 16]. Let us write a general instanton and antiinstanton solutions without vanishing fermionic classical coordinates [11, 13] in the following form

$$\begin{cases} q_+ = q_{0+} - \frac{\lambda_+ \lambda_-}{2V'(q_{0+})}, \psi_+ = \lambda_+ V'(q_{0+}), \psi_- = \frac{\lambda_-}{V'(q_{0+})}, \text{ inst.} \\ q_- = q_{0-} + \frac{\lambda_+ \lambda_-}{2V'(q_{0-})}, \psi_+ = \frac{\lambda_+}{V'(q_{0-})}, \psi_- = \lambda_- V'(q_{0-}), \text{ antiinst.} \end{cases} \quad (9)$$

where $\lambda_\pm$ are Grassmann integration constants and $q_{0\pm}(\tau)$ is a standard instanton/antiinstanton solution [16] which can be found from the equations

$$\dot{q}_{0\pm}(\tau) = \pm V'(q_{0\pm}). \quad (10)$$

The classical solutions (9) can be presented in more compact form

$$q_k = q_{0(k)} - \frac{\nu}{2W(q_{0(k)})}, \psi_{+(k)} = \lambda_+ W^k(q_{0(k)}), \psi_{-(k)} = \lambda_- W^{-k}(q_{0(k)}), \quad (11)$$

where $\nu = \lambda_+ \lambda_-$, $\nu^2 = 0$, $W(q) = V'(q)$, $k = +1$ for instantons and $k = -1$ for antiinstantons. The corresponding scalar superfields (1) are

$$\Phi_{(k)}(\tau, \theta^+, \theta^-) = q_{0(k)}(\tau) + \theta^+ \lambda_- W^{-k} + \theta^- \lambda_+ W^k + \theta^+ \theta^- \left(-W + \nu \frac{W'}{2W} \right). \quad (12)$$

It is important to stress that despite the fact that the classical trajectories have a nilpotent part proportional to ν (11) the action has no such additional parts and it is equal to the action on instanton trajectories, i.e.

$$S_{Ek} = \int W^2(q_{0(k)}) d\tau = S_{Eq_{0(k)}}. \quad (13)$$

For the topological charge Q_k of instantons/antiinstantons we obtain

$$Q_k = \frac{1}{2q_0(\infty)} \int_{-\infty}^{\infty} \dot{q}_k^2 d\tau = Q_{0(k)} + \frac{\nu}{2q_{(k)}(\infty)} \left[\frac{1}{W(q_{(k)}(+\infty))} - \frac{1}{W(q_{(k)}(-\infty))} \right]. \quad (14)$$

From (14) we conclude that the nilpotent part of the topological charge vanishes in case of spontaneous symmetry breaking, i.e. when $W(q) = V'(q)$ is an even function. That gives another meaning to the conclusions of [15, 16].

4 Fermionic Zero Modes

Let us consider a contribution of zero modes in quantum evolution of our system. In general we have

$$\Phi_{quant}(\tau, \theta^+, \theta^-) = \Phi(\tau + \tau_0, \theta^+ + \eta^+, \theta^- + \eta^-) + \Sigma', \quad (15)$$

where τ_0, η^+, η^- are shifts corresponding zero modes, Σ' is contribution of nonzero modes, $\Phi(\tau, \theta^+, \theta^-)$ is a classical superfield (12). Expanding the action next to the classical solution or differentiating of the classical equation of motion by corresponding parameter we find the equation for zero modes

$$\{1/2[D^+, D^-] - V''(\Phi)\}\Phi_0 = 0$$

But we can find the zero modes differentiating by the parameters of the solutions (9). We see that the zero modes connected to differentiating of ψ_- by λ_- and ψ_+ by λ_+ reduce to inverse functions. But from Cauchy-Schwarz-Bounjakovsky inequality

$$\left(\int_0^\infty \psi_{-0} dt\right)^2 \left(\int_0^\infty \psi_{+0} dt\right)^2 \leq \left(\int_0^\infty \psi_{-0} \times \psi_{+0} dt\right)^2$$

where ψ_{-0} and ψ_{+0} are fermionic zero modes, one can see that only one of two fermionic zero mode can be normalized and consequently one of two Grassmann constants have to vanish. But a vanished Grassmann constant corresponds to N=1 broken supersymmetry. Thus we see that one half of supersymmetries must be broken.

5 Supersymmetry as Anilpotence of Lagrangian

Let us consider nonsupersymmetric Lagrangian similar to (6) as

$$L_{E(nonsusy)} = \frac{\dot{q}^2}{2} + \frac{W^2(q)}{2} + \psi_+ \dot{\psi}_- + U(q) \psi_+ \psi_-, \quad (16)$$

where $W(q)$ and $U(q)$ are some functions. Equations of motion are

$$\ddot{q} - W(q) W'(q) + U'(q) \psi_+ \psi_- = 0, \quad (17)$$

$$\dot{\psi}_\pm \mp U(q) \psi_\pm = 0. \quad (18)$$

Fermionic classical solutions are

$$\psi_\pm = \lambda_\pm e^{\pm \int U(q) d\tau}, \quad (19)$$

and therefore $\psi_+ \psi_- = \lambda_+ \lambda_- = \nu = const$ on equations of motion. Let

$$q(\tau) = q_0(\tau) + \nu q_N(\tau), \quad (20)$$

where $q_0(\tau)$ is nonsupersymmetric instanton solution $\dot{q}_0 = \pm W(q_0)$ and $q_N(\tau)$ satisfies the following equation

$$\dot{q}_N = W'(q_0) q_N - \frac{U(q_0)}{W(q_0)}. \quad (21)$$

Then we obtain for on-shell Lagrangian (16)

$$L_{E(nonsusy)}^{n-shell} = W^2(q_0) + \nu W(q_0) [2W(q_0) W'(q_0) q_N - U(q_0)]. \quad (22)$$

The requirement of vanishing of the nilpotent part in (22) gives

$$q_N = \frac{U(q_0)}{2W(q_0) W'(q_0)}. \quad (23)$$

So that from (21) and (23) we finally obtain [11]

$$\frac{U'}{U} = \frac{W''}{W'} \quad (24)$$

which can be solved by

$$U = cW', \quad (25)$$

where c is an even constant which can be taken as 1. Substituting (25) into $L_{E(nonsusy)}$ (16) gives supersymmetric Lagrangian $L_{E(susy)}$ (cf. (6)) in the following form

$$L_{E(susy)} = \frac{\dot{q}^2}{2} + \frac{W^2(q)}{2} + \psi_+ \dot{\psi}_- + W'(q) \psi_+ \psi_-, \quad (26)$$

Thus, the requirement of anilpotence of the on-shell Lagrangian gives us the supersymmetry condition for the on-shell Lagrangian [11].

6 Quasiclassical Quantization Rules as Anilpotence of Action

Let us return to Minkowski (1|2) superspace by the substitution $\tau = it$. Then the action instead of (4) will take the form

$$S = \int L dt, \quad (27)$$

where³

$$L = \frac{\dot{q}^2}{2} - \frac{V'^2(q)}{2} + \psi_+ \dot{\psi}_- + V''(q) \psi_+ \psi_-. \quad (28)$$

The corresponding equations of motion are

$$\ddot{q} + V'(q) V''(q) - V'''(q) \psi_+ \psi_- = 0, \quad (29)$$

$$\dot{\psi}_\pm \pm iV''(q) \psi_\pm = 0. \quad (30)$$

Their solutions in case we admit nonzero fermionic classical solutions [13] have the following form

$$q = q_0 + \nu \frac{W}{2E}, \quad \psi_\pm = \lambda_\pm e^{\mp i \arcsin\left(\frac{W}{\sqrt{2E}}\right)}, \quad (31)$$

³In this section a dote denotes differentiation by t .

where E is energy of the system and $W = V'(q_0)$. Here $q_0(t)$ is a standard solution of 1-dimensional nonsupersymmetric system which can be found from the equation

$$dt = \frac{q_0}{\sqrt{2E - W^2}}. \quad (32)$$

The corresponding Maupertuis action is

$$S_M = \oint (pdq + \psi_+ d\psi_-), \quad (33)$$

where

$$p = \sqrt{2E - W^2} \left(1 + \nu \frac{W'}{2E} \right) \quad (34)$$

is a canonical momentum for q . On equations of motion the action (33) takes the form

$$S_M = \oint \sqrt{2E - W^2} dq_0 + \nu \oint \left(\frac{1}{E} - \frac{1}{2E - W^2} \right) \sqrt{2E - W^2} W' dq_0. \quad (35)$$

Second term with nilpotent addition vanishes after integration over a closed path and we obtain the standard super WKB rules [17].

The quasiclassical wave function is also defined by classical trajectories as follows

$$\Psi(q, \psi_+, \psi_-) \approx A_\psi^{\frac{i}{\hbar} S_\psi}, \quad (36)$$

where

$$A_\psi = \text{Ber} \left| -\frac{\partial^2 S_\psi}{\partial q(0) \partial q(t)} \right| \quad (37)$$

is a super generalization of the van Fleck determinant. Using the equations of motion (29)–(30) we obtain

$$A_\psi = \frac{1}{\sqrt{2E - W^2}} \left[1 + \frac{\nu}{2} \left(\frac{W'}{2E} + \frac{W'^2}{4E^2} e^{i \arcsin\left(\frac{W}{\sqrt{2E}}\right)} \right) \right], \quad (38)$$

$$S_\psi = \int dt (E - W^2) \left(1 + \nu \frac{W'}{E} \right). \quad (39)$$

Conclusions

Thus we have considered the generalized solutions for the classical equations of motion for a bosonic-fermionic system which contain the nilpotent additions – ”nilpotent marsh”. In consequence of conservation the nilpotent constant ν the particle will ”live” on such trajectory and never will have the intersections with usual classical trajectory. But on such instanton (antiinstanton) trajectories the classical action has a nilpotent addition, which vanishes only for supersymmetric Lagrangian. The nilpotent addition to the topological charge vanishes for the superpotential with spontaneous broken supersymmetry. We studied the normalized zero modes for the instanton and antiinstanton solutions. As a consequence of Cauchy-Schwarz-Bounjakovsky inequality for fermionic zero modes only half of them can be normalized. So we have to vanish half of Grassmann constants, which corresponds to $N = 1$ broken supersymmetry. In contrast of paper [4] we have used a superfield approach to SQM [14]. We have considered a quasiclassical action on such trajectories and obtained Super WKB rules for quantization, as an application of our approach.

Acknowledgments

We thank A. Pashnev, A. Burinsky, A. Gumenchuk and A. Zheltukhin for discussions and interest to the work.

This work is partly (A.V.) supported by grants INTAS-93/127ext, INTAS-93/493ext, INTAS-96/0308.

References

- [1] Witten E. *Dynamical breaking of supersymmetry* // Nucl. Phys. - 1981. - V. **B188**. - P. 513–537.
- [2] Andrianov A. A., Cannata F., Dedonder J.-P., Ioffe M. V. *SUSY quantum mechanics with complex superpotentials and real energy spectra*. - St. Petersburg: 1998. - 22 p. (Preprint / St. Petersburg State Univ.; SPbU-IP-97-24, quant-ph/9806019).
- [3] Diaconescu D.-E., Entin R. *A non-renormalization theorem for the $d = 1, N = 8$ vector multiplet* // Phys. Rev. - 1998. - V. **D56**. - P. 8045–8052.
- [4] Manton N. S. *Deconstructing supersymmetry*. - Cambridge: 1998. - 19 p. (Preprint / Univ. Cambridge; DAMTP 1998-39, hep-th/9806077).
- [5] Lowe D. A. *Eleven-dimensional Lorentz symmetry from SUSY quantum mechanics*. - Providence: 1998. - 16 p. (Preprint / Brown Univ.; BROWN-HET-1133, hep-th/9807229).
- [6] Paban S., Sethi S., Stern M. *Constraints from extended supersymmetry in quantum mechanics*. - Princeton: 1998. - 12 p. (Preprint / Inst. Adv. Study, hep-th/9805018).
- [7] Paban S., Sethi S., Stern M. *Supersymmetry and higher derivative terms in the effective action of Yang-Mills theories* // J. High Energy Phys. - 1998. - V. **06**. - P. 012–023.
- [8] Maldacena J. *The large N limit of superconformal field theories and supergravity*. - Cambridge: 1997. - 20 p. (Preprint / Harvard Univ.; HUTP-97/A097, hep-th/9711200).
- [9] Claus P., Derix M., Kallosh R., Kumar J., Townsend P. K., Van Proeyen A. *Black holes and superconformal mechanics*. - Stanford: 1998. - 9 p. (Preprint / Stanford Univ.; SU-ITP-98/27, hep-th/9804177).
- [10] de Wit B., Marquard U., Nicolai H. *Area preserving diffeomorphisms and supermembrane Lorentz invariance* // Comm. Math. Phys. - 1990. - V. **128**. - P. 39–56.

- [11] Akulov V. P. *New class of solutions for Boson-Fermion system of equations and supersymmetry* // *Ukrainian J. Phys.* - 1986. - V. **31**. - 11. - P. 1615–1618.
- [12] Akulov V. P., Volkov D. V. *Supersymmetric equations in the space (1, 2)* // *Theor. Math. Phys.* - 1980. - V. **42**. - P. 10–25.
- [13] Akulov V. P., Duplij S. *Quasiclassical quantization in supersymmetric quantum mechanics* // *Ukrainian J. Phys.* - 1988. - V. **33**. - 2. - P. 309–311.
- [14] Akulov V. P., Pashnev A. I. *Quantum superconformal model in (1, 2) space* // *Theor. Math. Phys.* - 1983. - V. **56**. - P. 862–878.
- [15] Cooper F., Freedman B. *Spontaneous supersymmetry breaking in quantum mechanics* // *Ann. Phys.* - 1983. - V. **146**. - 2. - P. 262–288.
- [16] Solomonson P., van Holten J. W. *Fermionic coordinates and supersymmetry in quantum mechanics* // *Nucl. Phys.* - 1982. - V. **B196**. - 3. - P. 509–531.
- [17] Comtet A., Bandrauk A. D., Campbell D. K. *Exactness of semiclassical bound states* // *Phys. Lett.* - 1985. - V. **B150**. - 1/3. - P. 159–162.